

PCT


WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁷ : A01N 43/16, A61K 31/35		A1	(11) International Publication Number: WO 00/40086
			(43) International Publication Date: 13 July 2000 (13.07.00)
(21) International Application Number: PCT/US00/00399 (22) International Filing Date: 7 January 2000 (07.01.00) (30) Priority Data: 09/226,580 7 January 1999 (07.01.99) US (71) Applicant: NITROSYSTEMS, INC. [US/US]; 512 Telfair Street, Augusta, GA 30901 (US). (72) Inventor: KAESEMEYER, Wayne, E.; 2433 McDowell Street, Augusta, GA 30904 (US). (74) Agent: MILLER, Raymond, A.; Reed Smith Shaw & McClay LLP, P.O. Box 488, Pittsburgh, PA 15230-0488 (US).		(81) Designated States: AE, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE, DK, DM, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, UZ, VN, YU, ZA, ZW, ARIPO patent (GH, GM, KE, LS, MW, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG). Published With international search report.	

(54) Title: FORMULATIONS FOR TREATING DISEASE AND METHODS OF USING SAME



(57) Abstract

A therapeutic mixture comprised of L-arginine and angiogenic growth factors is disclosed for the treatment of disease related to endothelial dysfunction. A schematic representation of NOS activation by select compounds and the conversion of L-arginine into NO is portrayed in the figure.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

FORMULATIONS FOR TREATING DISEASE AND METHODS OF USING SAME

5 This application claims continuing status under 35 U.S.C.
Section 120 from U.S. Patent Application Serial No. 08/833,842 filed April
10, 1997.

BACKGROUND OF THE INVENTION

10 This invention relates generally to therapeutic formulation
comprised of a combination of angiogenic growth factors and substrates of
Nitric Oxide Synthase ("NOS"), preferably arginine, as well as a method
of treating, preventing and/or ameliorating cardiocerebrorenovascular
disease and the symptoms thereof. More particularly, the present
invention is directed to the administration of a biological equivalent of
arginine and an agonist of NOS (*e.g.*, vascular endothelium growth factor
15 ("VEGF")) to produce a beneficial effect.

DESCRIPTION OF RELATED ART

20 One approach to treating cardiac disease is to effect the
dilation of vascular conduits in the body. In this regard, nitric oxide has
been shown to be formed enzymatically as a normal metabolite from
arginine in the vascular endothelium and provides an important
component to the formation of endothelium-derived relaxing factor
(EDRF). EDRF appears to be equivalent to Endothelium Derived Nitric

Oxide (EDNO) and as used herein EDRF and EDNO are used interchangeably unless otherwise indicated.

5 It has also been established that a family of enzymes called Nitric Oxide Synthase ("NOS") form nitric oxide from L-arginine, and the nitric oxide produced is responsible for the endothelium dependent relaxation and activation of soluble guanylate cyclase, neurotransmission in the central and peripheral nervous systems, and activated macrophage cytotoxicity. Nitric Oxide Synthase occurs in many distinct isoforms which include a constitutive form ("cNOS") and an inducible form (iNOS).
10 The constitutive form is present in normal endothelial cells, brain, neurons and some other tissues. Formation of nitric oxide by the constitutive form in endothelial cells is thought to play an important role in normal blood pressure regulation, prevention of endothelial dysfunction such as hyperlipodemia, arteriosclerosis, thrombosis, and restenosis. The
15 by-product of the conversion of L-arginine is L-citrulline. Brain, endothelium, and macrophage isoforms of NOS appear to be products of a variety of genes that have approximately 50% amino acid identity. NOS in brain and in endothelium have very similar properties, the major differences being that brain NOS is cytosolic and the endothelial enzyme
20 is mainly a membrane-associated protein.

Functionally, the constitutive form of Nitric Oxide Synthase, which is the predominant synthase present in brain and endothelium, may be active under basal conditions and can be further stimulated by increases in intracellular calcium that occur in response to receptor-mediated agonists or calcium ionophores. cNOS appears to be the
25 "physiological" form of the enzyme and plays a role in a diverse group of biologic processes. In vitro studies suggest that the activity of NOS can be

regulated in a negative feedback manner by nitric oxide itself. In
cardiocerebrorenovascular circulation, the primary target for
constitutively produced nitric oxide is believed to be soluble guanylate
cyclase located in vascular smooth muscle, the myocardium (myocytes)
5 and coronary vascular smooth muscle.

In contrast to the cNOS, the inducible, calcium-independent
form, iNOS was initially only described in macrophages. It is now known
that induction of nitric oxide synthase can occur in response to
appropriate stimuli in many other cell types. This includes both cells that
10 normally do not express a constitutive form of nitric oxide synthase, such
as vascular smooth muscle cells, as well as cells such as those of the
myocardium that express considerable levels of the constitutive isoform.
The inducible form of nitric oxide synthase has been found to be induced
in vascular smooth muscle cells, for example, by various cytokines and/or
15 microbial products.

iNOS exhibits negligible activity under basal conditions, but
in response to factors such as lipopolysaccharide and certain cytokines,
expression occurs over a period of hours. The induced form of the enzyme
produces much greater amounts of NO than the constitutive form, and
20 induced NOS appears to be the "pathophysiological" form of the enzyme
because high concentrations of NO produced by iNOS can be toxic to cells.
Induction of iNOS can be inhibited-by glucocorticoids and some cytokines.

SUMMARY OF THE INVENTION

The term "subject" as used herein to mean any mammal,
25 including humans, where nitric oxide formation from arginine occurs. The

methods herein for use on subjects contemplate prophylactic use as well as curative use.

5 The term endpoints as used herein refers to clinical events encountered in the course of treating cardiovascular disease, up to and including death (mortality).

10 L-arginine as used herein includes all biochemical equivalents (i.e. salts, precursors, and its basic form). "To mix", "mixing", or "mixture(s)" as used herein means mixing a substrate (i.e. L-arginine) and an agonist (i.e. angiogenic growth factors): 1) prior to administration ("in vitro mixing"); 2) mixing by simultaneous and/or consecutive, but separate (i.e. separate intravenous lines) administration of substrate (L-arginine) and agonist (angiogenic growth factor) to cause "in vivo mixing"; and 3) the administration of a NOS agonist after saturation with a NOS substrate (i.e. L-arginine is administered to build up a supply in the body prior to administering the NOS agonist (nitroglycerin or angiogenic growth factors)); or any combination of the above which results in the delivery of therapeutic amounts of a NOS agonist and a NOS substrate in an additive or synergistic way with regard to the treatment of disease, preferably cardiocerebrorenovascular disease.

20 Agonist refers to an agent which stimulates the bio-transformation of a substrate such as L-arginine either through enzymatic activation or increasing gene expression (i.e. increased levels of NOS). Of course, either or both of these mechanisms may be acting simultaneously.

The present invention is preferably useful in preventing, treating, arresting, or ameliorating disease conditions which are benefited by the bio-transformation of a substrate into nitric oxide or "native" nitric oxide. The present invention preferably promotes therapeutic angiogenesis.

The present invention may also be useful in preventing, treating, arresting, or ameliorating disease conditions which are benefited by the bio-transformation of L-arginine into "native" nitric oxide through enzyme activation of NOS.

The present invention may also be useful in achieving a beneficial effect when treating disease conditions by increasing or maximizing the production of EDRF or EDNO, and reducing clinical endpoints to include mortality.

The present invention may also be useful in preventing, treating, or avoiding tachycardia and ischemia.

The present invention may also be useful in preventing, treating, arresting, or ameliorating reperfusion injury in subjects who have had abrupt restoration of blood flow.

In one embodiment of the present invention, a mixture of NOS agonists, preferably angiogenic growth factors, and biological equivalents of L-arginine are used for the treatment of hypertension, hypertensive heart disease, coronary heart disease, including arteriosclerosis, angina, myocardial infarction, coronary thrombosis, restenosis post angioplasty, and sudden death, as well as a wide range of

cardiovascular disease (heart failure, stroke, and peripheral vascular diseases), and renovascular ischemia/hypertension.

In an alternative embodiment of the invention, therapeutically effective amounts of a precursor of EDNO and an agonist of NOS (preferably one of those cited in Table I herein) are combined prior to administration to a patient. In another embodiment of the invention, therapeutically effective amounts of a precursor of EDNO and an angiogenic growth factor are combined and administered.

In a preferred embodiment of the present invention, therapeutically effective amounts of L-arginine and therapeutically effective amounts of a macrophage secretory product are mixed at a physiologically acceptable pH.

Another embodiment of the present invention is a method for treating hypertension in a subject by vasodilation or vasorelaxation comprising: selecting a hypertensive subject; administering L-arginine and a NOS agonist (*e.g.*, those listed in Table I), preferably angiogenic growth factors, to the subject; obtaining periodic blood pressure measurements of the subject; and continuing administration of L-arginine and the NOS agonist, preferably a secretory product, more preferably an angiogenic growth factor, even more preferably VEGF or bFGF angiogenic growth factors until a desirable blood pressure or therapeutic effect is detected in the subject. A desirable blood pressure in a hypertensive subject should ultimately be within the following ranges: systolic preferably in the range of 95-180 mmHg, more preferably in the range of 105-165 mmHg, and even more preferably in the range of 120 to 140 mmHg; and diastolic preferably in the range of 55-115 mmHg, more

preferably in the range of 65-100 mmHg, and even more preferably in the range of 70 to 90 mmHg, and most preferably 75-85 mmHg. Under no circumstances should the systolic be permitted to go below 95 mmHg.

Another embodiment of the present invention is a method for
5 preventing or treating cardiovascular disease in a non-hypertensive
subject by vasodilation or vasorelaxation comprising: selecting a subject;
administering to said subject a formulation comprising a mixture of an
angiogenic growth factor and an endothelium dependent source or
precursor of nitric oxide (*e.g.*, L-arginine); obtaining periodic
10 measurements of vasorelaxation on the subject and; continuing
administration of the formulation until a desirable state of vasorelaxation
or desirable therapeutic (*e.g.*, angiogenic) effect is detected in the subject.

Yet another embodiment is a method for stimulating cNOS
in a subject which comprises: selecting a subject; administering to said
15 subject a formulation comprising a mixture of L-arginine and angiogenic
growth factors so as to increase "native" NO production and reduce
endpoints to include mortality.

BRIEF DESCRIPTION OF THE DRAWINGS

Fig. 1 is a schematic representation of NOS activation by
20 select compounds and the conversion of L-arginine into NO.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Research into the area of NOS activation reveals a number of agonists of NOS. Interestingly, many of the NOS agonists have also been implicated in angiogenesis. Substance P ("SP"), a secretory product, is identified herein as a cNOS agonist. Other secretory products (e.g., those identified in "Macrophages and angiogenesis" by Sunderkotter et al. (J Leukoc Biol 1994 Mar; 55(3):410-22)) may also be expected to be agonists of NOS. Bradykinin ("BK"), a NOS agonist, has also been implicated as a possible angiogenic factor. Angiogenic growth factors like those identified in Table I stimulate the growth of new blood vessels (e.g., in vascular beds such as the coronary, peripheral, etc.) previously occluded with atherosclerotic obstructions. Angiogenic growth factors are proteins which were initially discovered as agents responsible for the growth of new blood vessels which maintain the growth and spread of cancerous tumors (neovascularization). Two of the angiogenic growth factors, vascular endothelial growth factor (VEGF) and basic fibroblastic growth factor (bFGF), have been infused into catheters, used at the time of generating coronary and peripheral arteriograms, and have resulted in the growth of significant new collateral blood vessels in the region of ischemia producing vascular atherosclerotic occlusions. In this way, the symptoms of ischemia are lessened. The term applied to this treatment approach is "therapeutic angiogenesis."

Like angiogenic agents Substance P and Bradykinin, VEGF and bFGF also appear to act as NOS agonists, specifically cNOS. It appears the resultant EDNO produced is in large part responsible for the new collateral vessel growth ("collateral") which in turn is responsible for

the improvement in symptoms of ischemia seen in therapeutic angiogenesis. Furthermore, it has also been shown that the collateral responses to both VEGF and bFGF can be magnified significantly with L-arginine supplementation. Therefore, angiogenic growth factors, preferably VEGF and bFGF, appear to have dual applicability in the treatment of hypertension and cardiovascular diseases in that they both stimulate therapeutic angiogenesis and activity of Nitric Oxide Synthase. It also appears that the overall therapeutic angiogenic result with angiogenic growth factors is augmented to the extent they act as agonists of NOS. The fact that angiogenic growth factors are agonists or stimulators of nitric oxide synthase has important implications. Mixing angiogenic growth factors "in vitro" or "in vivo" with L-arginine may have an unforeseen beneficial effect that is associated with excess L-arginine providing additional substrate for NOS and the NOS being catalyzed to enzymatically increase the bio-transformation of L-arginine into nitric oxide (EDRF or EDNO) which would in turn amplify the overall therapeutic effect.

Stimulation of NOS by angiogenic growth factor(s) in the presence of excess L-arginine or other substrate precursor of native NO may be used to prevent, treat, arrest, or ameliorate any disease or condition which is positively affected by NO production. Such conditions include hypertensive cardiocerebrorenovascular diseases and their symptoms as well as non-hypertensive cardiocerebrorenovascular diseases. The mixture is particularly useful for subjects in need of native NO production for therapeutic angiogenesis. Application of such a mixture is beneficial for: (1) chronic stable angina; (2) unstable angina; (3) acute myocardial infarction; (4) hibernating myocardium; (5) stunned

myocardium; (6) limitation of ventricular remodeling in post myocardial infarction and subsequent risk of congestive heart failure; (7) prophylaxis of recurrent myocardial infarction; (8) prevention of sudden death following myocardial infarction; (9) vasospastic angina; (10) congestive heart failure-systolic-seen in association with 1-6 above; (11) congestive heart failure-diastolic-seen in association with 1-10 above and 12-15 below; (12) microvascular angina seen in association with 1-11 above and 15 and 16 below; (13) silent ischemia seen in association with 1-12 above and 15 and 16 below; (14) reduction of ventricular ectopic activity seen in association with 1-13 above and 15 below; (15) any or all of the above 1-14 states of ischemic myocardium associated with hypertensive heart disease and impaired coronary vasodilator reserve; (16) control of blood pressure in the treatment of hypertensive crisis, perioperative hypertension, uncomplicated essential hypertension and secondary hypertension; (17) regression of left ventricular hypertrophy seen in association with 15 and 16 above; (18) prevention and or regression of epicardial coronary arteriosclerosis seen in 1-17 above; (19) prevention of restenosis post angioplasty; (20) prevention and/or amelioration of free radical mediated reperfusion injury in association with 1-19 above; (21) use of the combination in the prevention of myocardial injury during cardioplegic arrest during coronary bypass or other open heart surgery i.e. use of the combination as a cardioplegic solution; (22) post transplant cardiomyopathy; (23) renovascular ischemia; (24) cerebrovascular ischemia (TIA) and stroke); (25) pulmonary hypertension; and (26) peripheral vascular disease (claudication).

Vascular smooth muscle cells are located mainly in veins, arteries, and coronary arteries. The following discussion focuses on

smooth muscle and myocyte relaxation stimulated by vasodilators but should not be so limited. The present invention is useful when NO regulation is beneficial. As discussed above the Nitric Oxide Synthase in the cells is normally cNOS, the constitutive form of Nitric Oxide Synthase, and the generator cells are endothelial cells and the target cells are vascular smooth muscle cells. Fig. 1 is a schematic illustration of a proposed mechanism of action of preferred substances (e.g., angiogenic growth factors) and arginine and is not intended to imply any cellular relationship or geography of the various sites of action, but rather meant to illustrate their functional relationship. Although Fig. 1 list certain preferred agents as angiogenic agents, it is meant as a representative sampling of those listed in Table I. In Fig. 1 the abbreviations corresponding to Table I are used with SP representing Substance P and GF representing select Growth Factors.

The principle combination to be employed will be a mixture that involves therapeutic concentrations of L-arginine and therapeutic concentrations of angiogenic growth factor in water. Any pharmaceutical grade L-arginine will be sufficient and should be diluted preferably to 2.5-60% w/v (g/ml), more preferably to 5-45% w/v (g/ml), even more preferably between 7.5-30% w/v (g/ml), even more preferably to 10-15% w/v (g/ml), and most preferably 10% w/v (g/ml) L-arginine. The typical doses anticipated will be 30 grams of L-arginine in sterile water (Total Volume 300 cc). L-arginine is anticipated eventually to be approximately 10:1 to about 25:1 of the hydrochloride salt to L-arginine as a base, and even more preferably 15:1 to about 20:1 hydrochloride salt to base, and most preferably 15:1 hydrochloride salt to base. In this example 28 to 29 grams will be the hydrochloride salt and 1 to 2 grams of L-arginine will be base.

In the preferred embodiment discussed herein, L-arginine is used in conjunction with any of the family of those substances known as angiogenic growth factors. However, those particular angiogenic growth factors most preferred for use in conjunction with the present formulation are selected from the group consisting of VEGF and bFGF and even more preferably VEGF. Any of the agonists of Table I may be suitable candidates for use in combination with L-arginine.

As part of a "mixture", the angiogenic growth factor is included together with L-arginine and clinically effective weight ratios of between 1:2 to 1:150. Even more particularly, the ratio of the angiogenic growth factor to L-arginine in the formulation is between 1:5 to 1:100. The most preferred embodiment of the "mixture" the ratio of angiogenic growth factor, more preferably VEGF or bFGF, to L-arginine is 1:50. VEGF can be obtained from Genentech (South San Francisco, CA) and bFGF can be obtained from R&D Systems (Minneapolis, MN). The range of ratios of an angiogenic growth factor to L-arginine may be employed with virtually any of the angiogenic growth factors.

Where the particular angiogenic growth factor is VEGF the ratio of VEGF to L-arginine is preferably within the range 1:2 to 1:50, Wt/Wt. For example, VEGF/L-arginine at a ratio of 1:2 would include 40 mg/day VEGF with 80 mg/day L-arginine. Where the ratio of VEGF/ L-arginine is at a ratio of 1:20, for example, 20 mg/day VEGF would be administered with 400 mg/day L-arginine. Weight ratio of ingredients described herein in regard to VEGF or bFGF are applicable for any of the agents listed in Table I. The amounts above have been found to be effective, however, each route of administration (i.e. IV, oral, transdermal, intracoronary, intra-arterial, etc.) may vary in their requirements.

Even more particularly, the presently disclosed "mixtures" may be described in terms of their relative concentrations (grams) administered as part of a continuous intracoronary, intra-arterial, intravenous and intrapericardial infusions. In one particular embodiment, the formulation is administered as mixtures of growth factors with L-arginine encased in liposomes so as to provide maximum retention time of the mixture in any given vascular bed being perfused by a catheter delivering the growth factor/ L-arginine angiogenic mixture. In some cases the liposomes containing the mixture of growth factors and L-arginine may also contain genetic material which will code for the synthesis of the growth factor following transfection of the genetic material into the surrounding tissue of the vascular bed. In some cases pellets containing the aforementioned mixtures may be directly implanted into the myocardium at the time of coronary bypass graft surgery. In yet another case, a therapeutic mixture of L-arginine and an angiogenic growth factor are repeatedly infused into the pericardial space via an indwelling infusion catheter.

By way of example only, Table I presents a listing of several agents, most of which are angiogenic growth factors. These substances vary in their potency and their abilities to activate nitric oxide synthase and stimulate therapeutic angiogenesis.

TABLE I

EGF (epidermal growth factor)	TGF- α (transforming growth factor- α)	Substance P
FGF-1 (fibroblast growth factor-1; aFGF)	TGF- β (transforming growth factor- β)	TAT (HIV-1 transactivating protein)
FGF-2 (fibroblast growth factor-2; bFGF)	TNF- α (tumor necrosis factor- α)	UPA (urokinase)
FGF-3 (fibroblast growth factor-3; int-2)	TP (thymidine phosphorylase; also called platelet-derived endothelial cell growth factor [PD-ECGF])	VCAM-1 (vascular cell adhesion molecule)
FGF-4 (fibroblast growth factor-4; hst-1, K-FGF)	VEGF (vascular endothelial growth factor; vascular permeability factor [VPF]; vasculotropin)	Lactate
Bradykinin (BK)	Angiogenin	Hyaluronan fragments
G-CSF (granulocyte colony-stimulating factor)	Angiotensin II	Erucamide
IGF-1 (insulin-like growth factor-1)	Ceruloplasmin	Leukotrienes C ₄ and D ₄
IL-8 (interleukin-8)	Polyamines	PGE ₁ , PGE ₂ , (prostaglandins E ₁ , E ₂)
PDGF (platelet-derived growth factor)	Platelet activating factor	Estrogen
SF (scatter factor; same as hepatocyte growth factor, [HGH])	Proliferin	Nicotinamide

Also specifically within the scope of those agents listed in Table I are their bio-active metabolites. Any one or several of those angiogenic growth factor compounds listed in Table 1 may be mixed with L-arginine or substrate precursor to endogenous nitric oxide to provide a therapeutically effective treatment for a patient.

While not wishing to be bound by theory, it is now believed that angiogenic agents have a stimulating effect on cNOS, and furthermore, that this action to stimulate cNOS is involved in the angiogenic response seen with growth factors. Thus, angiogenic growth factors and L-arginine appear to have a heretofore unexpected additive and/or synergistic effect on cNOS stimulation. The stimulation of cNOS may be a result of cNOS having a unique receptor site for angiogenic growth factors or angiogenic growth factors may initiate a cascade of events which stimulate NOS. Administering the two also provides adequate substrate for cNOS processing of L-arginine since the L-arginine is added in excess while at the same time stimulation the enzymatic activity of NOS. Whether it is a synergistic effect or additive effect, what is clear is that "mixing" a precursor substrate of "native" nitric oxide with angiogenic agents (*e.g.*, VEGF and bFGF) results in a heretofore unexpected increase in NO production and angiogenesis. While not wishing to be bound by theory, growth factors appear to stimulate cNOS in much the same way as other NOS agonist and cNOS stimulation in turn stimulates angiogenesis.

By way of example, a therapeutic mixture, as previously described, of L-arginine and VEGF is infused into the coronary arteries of a patient with chest pain at the time of a coronary arteriogram which

reveals multiple diffuse atherosclerotic obstructive lesions which are amenable to treatment with bypass surgery or angioplasty. The treatment results in improvement in collateral blood flow and a decrease in the severity and frequency of angina attacks over the next 6 to 12 months.

The methods of the present invention involve administering to a mammalian host, preferably a human host, pharmacologically effective amounts of arginine and angiogenic growth factor. The agents may be combined in vitro before administration or separately administered, either concurrently or simultaneously, with administration generally taking place up to 24 hours before or after the administration of the other biological active agent(s).

The administration(s) may take place by any suitable technique, including oral, subcutaneous and parenteral administration, preferably parenteral or oral. Examples of parenteral administration include intravenous, intra-arterial, intramuscular, and intraperitoneal. The dose and dosage regimen will depend mainly on whether the inhibitors are being administered for therapeutic or prophylactic purposes, separately or as a mixture, the type of biological damage and host, the history of the host, and the type of inhibitors or biologically active agent. The amount must be effective to achieve an enhanced therapeutic index. It is noted that humans are generally treated longer than the mice and rats with a length proportional to the length of the disease process and drug effectiveness. The doses may be single doses or multiple doses over a period of several days. Therapeutic purposes is achieved as defined herein is when the treated hosts exhibit improvement against disease or infection, including but not limited to improved

survival rate, more rapid recovery, or improvement or elimination of symptoms. If multiple doses are employed, as preferred, the frequency of administration will depend, for example, on the type of host and type of cancer, dosage amounts, etc. The practitioner may need to ascertain upon
5 routine experimentation which route of administration and frequency of administration are most effective in any particular case.

Compounds and agents (e.g., both L-arginine and angiogenic growth factors) of the present invention, in conjunction with a pharmaceutically acceptable carrier, may be used for any of the
10 therapeutic effects, discussed above. Such compositions may be in the form of an agent(s) in combination with at least one other agent, such as stabilizing compound, which may be administered in any sterile, bio-compatible pharmaceutical carrier, including, but not limited to, saline, buffered saline, dextrose, and water. The compositions may be
15 administered to a patient alone, or in combination with other agents, drugs or hormones. Pharmaceutically-acceptable carriers may also be comprised of excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. Further details on techniques for formulation and administration may be
20 found in the latest edition of Remington's Pharmaceutical Sciences (Maack Publishing Co., Easton, PA) hereby incorporated herein by reference in its entirety.

Pharmaceutical compositions for oral administration can be formulated using pharmaceutically acceptable carriers well known in the
25 art in dosages suitable for oral administration. Such carriers enable the pharmaceutical compositions to be formulated as tablets, pills, dragees,

capsules, liquids, gels, syrups, slurries, suspensions, and the like, for ingestion by the patient.

Pharmaceutical preparations for oral use can be obtained through combination of active compounds with solid excipient, optionally grinding a resulting mixture, and processing the mixture of granules, after adding suitable auxiliaries, if desired, to obtain tablets or dragee cores. Suitable excipients are carbohydrate or protein fillers, such as sugars, including lactose, sucrose, mannitol, or sorbitol; starch from corn, wheat, rice, potato, or other plants; cellulose, such as methyl cellulose, hydroxypropylmethyl-cellulose, or sodium carboxymethylcellulose; gums including arabic and tragacanth; and proteins such as gelatin and collagen. If desired, disintegrating or solubilizing agents may be added, such as the cross-linked polyvinyl pyrrolidone, agar, alginic acid, or a salt thereof, such as sodium alginate.

Dragee cores may be used in conjunction with suitable coatings, such as concentrated sugar solutions, which may also contain gum arabic, talc, polyvinylpyrrolidone, carbopol gel, polyethylene glycol, and/or titanium dioxide, lacquer solutions, and suitable organic solvents or solvent mixtures. Dyestuffs or pigments may be added to the tablets or dragee coatings for product identification or to characterize the quantity of active compound. i.e., dosage.

Pharmaceutical preparations which can be used orally include push-fit capsules made of gelatin, as well as soft, sealed capsules made of gelatin and a coating, such as glycerol or sorbitol. Push-fit capsules can contain active ingredients mixed with a filler or binders, such as lactose or starches, lubricants, such as talc or magnesium

stearate, and, optionally, stabilizers. In soft capsules, the active compounds may be dissolved or suspended in suitable liquids, such as fatty oils, liquid, or liquid polyethylene glycol with or without stabilizers.

Pharmaceutical formulations suitable for parenteral administration may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiologically buffered saline. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran.

Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Optionally, the suspension may also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

For topical or nasal administration, penetrants appropriate to the particular barrier to be permeated are used in the formulation. Such penetrants are generally known in the art.

The pharmaceutical compositions of the present invention may be manufactured in a manner that is known in the art, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levigating, emulsifying, encapsulating, entrapping, or lyophilizing processes.

The pharmaceutical composition may be provided as a salt and can be formed with many acids, including but not limited to, hydrochloric, sulfuric, acetic, lactic, tartaric, malic, succinic, etc. Salts tend to be more soluble in aqueous or other protonic solvents than are the corresponding free base forms. In other cases, the preferred preparation may be a lyophilized powder which may contain any or all of the following: 1-50 mM histidine, 0.1%-2% sucrose, and 2-7% mannitol, at a pH range of 4.5 to 5.5, that is combined with buffer prior to use.

After pharmaceutical compositions have been prepared, they can be placed in an appropriate container and labeled for treatment of an indicated condition. Such labeling would include amount, frequency, and method of administration.

The therapeutically effective dose of specific angiogenic growth factors can be estimated initially either in cell culture assays, e.g., of neoplastic cells, or in animal models, usually mice, rabbits, dogs, or pigs. The animal model may also be used to determine the appropriate concentration range and route of administration. Such information can then be used to determine useful doses and routes for administration in humans.

The exact dosage will be determined by the practitioner, in light of factors related to the subject that requires treatment. Dosage and administration are adjusted to provide sufficient levels of the active moiety or to maintain the desired effect. Factors which may be taken into account include the severity of the disease state, general health of the subject, age, weight, and gender of the subject, diet, time and frequency of administration, drug combination(s), reaction sensitivities, and

tolerance/response to therapy. Long-acting pharmaceutical compositions may be administered every 3 to 4 days, every week, or once every two weeks depending on half-life and clearance rate of the particular formulation.

5 Although the preferred methods have been described in detail, it should be understood that various changes, substitutions, and alterations can be made in the present invention as defined by the claims appended hereto. For example, the aforementioned therapeutic mixture of L-arginine and VEGF may alternatively be infused into a catheter used
10 to diagnose peripheral vascular disease in patients suffering from claudication. The present invention is to be defined only by the claims attached hereto.

What Is Claimed Is:

1. A therapeutic mixture of an angiogenic growth factor and a substrate of NOS.
2. The therapeutic mixture of claim 1, which said
5 substrate of NOS is a biological equivalent of L-arginine.
3. The therapeutic mixture of claim 1, wherein said substrate of NOS is L-arginine.
4. The therapeutic mixture of claim 1, wherein the
10 angiogenic growth factor is selected from the group consisting of: EGF, TGF- α , TGF- β , FGF-1, FGF-2, FGF-3, FGF-4, VEGF, and IGF-1.
5. The therapeutic mixture of claim 1, wherein said angiogenic growth factor is selected from the group consisting of:

VEGF; and

bFGF.
- 15 6. A method of treating a disease condition in a subject by vasodilation or vasorelaxation comprising:

selecting a subject;

administering a mixture of L-arginine and an angiogenic growth factor

obtaining periodic indicators of vasorelaxations for the
subject; and

continuing administration of the mixture until a desirable
state of vasorelaxation is obtained.

5 7. The method of claim 6, wherein the formulation is
administered intravenously, buccal, intracoronary, intraarterially,
intramuscularly, topically, intranasally, rectally, sublingually, orally,
subcutaneously, by patch, or inhalation.

10 8. The method of claim 6, wherein said disease is
hypertension, hypertensive heart disease, coronary heart disease,
cardiovascular disease, cerebrovascular disease, and renovascular disease.

9. The method of claim 8, wherein said coronary heart
disease is restenosis post angioplasty.

15 10. The method of claim 6, wherein L-arginine and said
angiogenic growth factor are mixed in vivo.

11. The method of claim 10, wherein L-arginine and said
angiogenic growth factor are administered at a therapeutic concentration.

12. The method of claim 11, wherein the therapeutic
concentration of L-arginine is from 7.5% to about 30% w/v (g/ml).

13. The method of claim 6, wherein said angiogenic growth factor is selected from the group consisting of EGF, TGF- α , TGF- β , FGF-1, FGF-2, FGF-3, FGF-4, VEGF, and IGF-1.

5 14. The method of claim 6, wherein said angiogenic growth factor is selected from the group consisting of VEGF and bFGF.

15. The method of claim 6, wherein said angiogenic growth factor is VEGF.

16. The method of claim 6, wherein said angiogenic growth factor is bFGF.

10 17. A method of stimulating nitric oxide synthase to produce nitric oxide, said method comprising:

administering L-arginine and an agonist of nitric oxide synthase to a subject have a nitric oxide synthase receptor site, said agonist being different than L-arginine and being an angiogenic growth factor; and
15

stimulating said nitric oxide synthase to a desirable level with said agonist of nitric oxide synthase.


18. The method of claim 17, wherein said L-arginine is in excess of said angiogenic growth factor.

20 19. The method of claim 17, wherein therapeutically effective amounts of L-arginine is combined with therapeutically effective

amounts of said angiogenic growth factor prior to administering to the patient.

20. The method of claim 17, wherein angiogenic growth factor is selected from the group consisting of EGF, TGF- α , TGF- β , FGF-1, FGF-2, FGF-3, FGF-4, VEGF, and IGF-1.
- 5

Fig. 1



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US00/00399

A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : A01N 43/16; A61K 31/35

US CL : 514/460

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/460

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched
NONEElectronic data base consulted during the international search (name of data base and, where practicable, search terms used)
NONE

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	SUNDERKOTTER et al. Macrophages and angiogenesis. J. Leukoc. Biol. March 1994, Vol. 55, NO. 3, pages 410-420, see entire document.	1-20
Y	BASSENGE, E. Coronary Vasomotor Responses: Role of Endothelium and Nitrovasodilators. Cardiovascular Drugs and Therapy. 1994, Vol. 8, pages 601-610, see entire document.	1-20



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
E earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*-&* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

16 MARCH 2000

Date of mailing of the international search report

18 APR 2000

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

DWAYNE C. JONES

Telephone No. (703) 308-1235